




STEPHEN R. SHOROFKY, M.D., PH.D.
Director
MAGDI SABA, M.B., B. Ch., M.Sc.
R.E. HOOD, M.D.

ROBERT W. PETERS, M.D.
TIMM L. DICKFELD, M.D., PH.D.
ANASTASIOS P. SALIARIS, M.D.

UNIVERSITY OF MARYLAND PHYSICIANS, P.A.

LOVENOX INJECTIONS



Injections are given in places on your body where you have fatty tissue. The best sites are: abdomen (stomach), back of the arm, top of leg, and hip area.

TO INJECT:

1. Prepare the injection site by cleaning the area with an alcohol swab. To do this start at the center, apply pressure, and cleanse in a circular motion working outwards. Do not retrace your steps.
2. Wait a few seconds until the alcohol has dried. This reduces the sting. Remove the needle guard from the needle and syringe filled medication.
3. With your freehand, gently pinch up the skin at the injection site. Hold the syringe like a pencil and push the needle straight into the skin.
4. When the needle is in place, slowly pull back on the plunger to see if any blood flows into the syringe. If some blood does enter the syringe (a rare occurrence), remove the needle, discard the syringe and the medication, and prepare another injection.
5. If no blood enters the syringe, slowly inject the medication by gently pushing the plunger down with your index finger until the syringe is empty.
6. Release the pinch, and remove the needle quickly. Apply pressure to the injection site with a dry, sterile gauze. You may notice a drop of blood, but there is no cause for concern. If desired, apply a Band-Aid
7. Discard your used syringe into an opaque, puncture proof non-breakable container.

Revised
04/06